

Id-dijanjosi tad-dimenzja

MALTESE | ENGLISH

Dan il-Fuljett ta' Ghajnuna jagħti tagħrif dwar is-sinjalibikrin tad-dimenzja, t-teknika użata biex issir id-dijanjosi tad-dimenzja u l-importanza ta'dijanjosi bikrija u korretta.

X'inħuma s-sinjalibikrin tad-dimenzja?

Billi il-marda li tikkawża d-dimenzja tiżviluppa bil-mod is-sinjalibikrin jistgħu jkunu sotili ħafna u ma jkunux jidher malajr. Is-sintomi bikrin jiddependu wkoll mit-tip ta-dimenzja u jvarjaw ħafna minn persuna għall-oħra.

Sintomi bikrin komuni jinkludu:

- Problemi tal-memorja speċjalment biex jiftakru ġrajjet riċenti
- Konfużjoni li tiżdied
- Tnaqqis fil-konċentrazzjoni
- Tibdin fil-personalità jew fl-imġiba
- Apatija u rtirar jew dipressjoni
- Telfien tal-kapaċită biex jagħmlu xogħlijiet ta' kuljum

Kultant in-nies ma jintebħux li dawn is-sintomi jindikaw li xi ħaġa mhijiex f'posta. Jistgħu b'mod żbaljat jassumu li mīġiba bħal dik hija parti normali tal-proċess tax-xjuhiha. Inkella s-sintomi jistgħu jiżviluppaw gradwalment u ma jkunux innotati għal żmien twil. Kultant xi wħud ma jkunux iridu jaġixxu wkoll meta jkunu jafu li xi ħaġa mhijiex sewwa.

Għal persuna li jkollha dawn is-sintomi, n-natura stess tat-tibdiliet fil-moħħiġ tista' tħisser li l-persuna mhiex kapaci tirrealizza li hemm it-tibdil.

Sinjalibikrin ta' twissija

Din hija lista biex tiċċekkja s-sintomi komuni tad-dimenzja. Imxi mal-lista u mmarka s-sintomi li jeżistu. Jekk tniżżeż ħafna marki, kellem tabib biex jassessjak sewwa.

Telfien tal-memorja li jaffettwa l-funzjonament ta' kuljum

Huwa normali li kultant tinsa l-appuntamenti u tiftakarhom aktar l-quddiem. Persuna bid-dimenzja tista' tħalli l-affarijiet iktar ta'sikwit jew ma tiftakarhom qatt.

Diffikultà biex tagħmel xogħlijiet familjari.

In-nies jistgħu jkunu distratti u jistgħu jinsew iservu parti mill-ikla. Persuna bid-dimenzja tista' jkollha diffikultà bil-passi kollha meħtieġa biex tħejji ikla.

Konfużjoni fil-ħin u l-post

Persuna bid-dimenzja jista' jkollha diffikultà biex issib t-triq lejn post familjari jew tkun konfużja dwar fejn qiegħda jew taħseb li qiegħda lura fi żmien mgħoddha ta' ħajjitha.

Problemi bil-lingwa

Kultant kulħadd ikollu diffikultà biex isib il-kelma eżatta, iżda persuna bid-dimenzja tista' tħalli tħafif jew iddaħħal minflokkom kliem li ma joqgħodx u jkun diffiċċi biex tifhimhom. Jista' jkun ukoll li jsibu diffiċċi li jifhmu lill-oħrajn.

Problemi bil-ħsebijiet astratti

L-immaniġġar tal-finanzi jista' jkun tqil għal kulħadd iżda persuna bid-dimenzja tista' ssibha bi tqila biex tkun taf xi jfissru n-numri u x'tagħmel bihom.

Ġudizzju fqir jew nieqes

Ħafna attivitajiet jinħtieġu ġjudizzju tajjeb. Meta din il-kapiċċità tkun affewwa mid-dimenzja l-persuna jista' jkollha diffikultà biex tagħmel deċiżjonijiet xierqa, bħal x'tilbes f'temp kiesaħ.

Problemi bil-kapaċitajiet spazjali

Persuna bid-dimenzja ssibha diffiċċi biex tikkalkula d-distanza jew id-direzzjoni meta tkun qiegħda ssuq.

Problem bit-telf tal-ogġetti

Kulħadd jista' jitlef għal fit il-kartiera jew iċ-ċwievet. Persuna bid-dimenzja tista' tqiegħed spiss l-ogġetti f'postijiet mhux addattati.

Bidliet fil-buri, fil-personalità jew l-imġiba

Kulħadd ikun imdejjaq jew bil-buri minn żmien għal żmien. Dawk bid-dimenzja jistgħu jibdu l-buri malajr għall-ebda raġuni apparenti. Jistgħu jsiru konfużi, suspettużi jew irtrati fihom nfushom. Xi wħud jistgħu ma jibqghux jistħu jew isiru jitkellmu iktar.

Telfien tal-inizjattiva

Huwa normali li tiddejjaq minn xi attivitajiet. Madankollu d-dimenzja tista' ġġiegħel persuna titlef l-interess f'attivitajiet li qabel kienet tieħu gost tagħmel.

National Dementia Helpline **1800 100 500**

dementia.org.au

Tista' ma tkunx dimenžja

Fitakar li ħafna kundizzjonijiet għandhom sintomi jixbhu lil dawk tad-dimenzja, għalhekk tassumix li xi ħadd ikollu d-dimenzja għax ikollu xi ffit mis-sintomi msemmija hawn fuq. Puplesji, d-dipressjoni, l-alkoholiżmu, l-infezzjonijiet, id-diżordnijiet tal-ormoni, nuqqas ta'nutrizzjoni u t-tumuri tal-moħħi jistgħu kollha jkollhom sintomi bħal tad-dimenzja. Hafna minn dawn il-kundizzjonijiet jistgħu jkunu kukturati.

Dijanjosi korretta hija importanti

Huwa importanti li t-tabib ikun kkonsultat fi stadju bikri. Tabib biss jista' jagħmel dijanjosi tad-dimenzja. Assessjar mediku kompli jista' jidher kundizzjoni li tista' tkun ikkurata u jiġura li tkun ikkurata b'mod korrett jew jikkonferma il-preżenza tad-dimenzja.

Assessjar jista' jinkludi dan li ġej:

- Storja medika – it-tabib se jistaqsi dwar il-problemi medikalji tal-imgħoddxi u tal-preżent, l-istorja medika tal-familja, x'mediciċi li qed jittieħdu, u l-problemi bil-memoria, bil-ħsieb jew bl-imġiba li jkunu jikkawżaw thassib. It-tabib jista' jkun irid ukoll ikellem lil xi membru qarib tal-familja li jkun jista' jgħin biex tkun provduta l-informazzjoni neċċessarja kollha.
- Ezami fiziku – dan jista' jkun jinkludi testijiet tas-sensi u l-funzjonament tal-movimenti, kif ukoll il-funzjoni tal-qalb u l-pulmun biex jgħinu ħalli jkunu eskużi kundizzjonijiet oħra.
- Testijiet fil-laboratorju – li jinkludu varjetà ta' testijiet tad-demm u l-awrina biex tkun identifikata l-possibilita` ta' marda li tkun responsabbi għas-sintomi. F'xi kazijiet jista' jkun meħud kampjun żgħir tal-fluwidu mis-sinsla biex ikun it-testejt.
- Test newropsikoloġiku jew konjittiv – jintużaw varjetà ta' testijiet biex ikunu assessjati ll-kapaċitajiet tal-ħsieb inkluzi l-memoria, l-lingwa, l-attenzjoni u s-soluzzjoni tal-problemi. Dan jista' jgħin biex ikunu identifikati iz-zoni ta' problemi specifiċi, li mbagħad jgħin biex jiddentifikasi l-kawża bażika jew t-tip ta' dimenžja.
- Immaġni tal-moħħi – hemm certi skannijiet li jaraw l-istruttura tal-moħħi u li huma wżati biex jaqtgħu barra t-tumuri tal-moħħi jew demm magħqud fil-moħħi bħala raġuni għas-sintomi, u biex isibu in-nisġa tat-telf tat-tessuti tal-moħħi li tista' jagħmel differenza bejn it-tipi differenti ta' dimenžja. Skannijiet oħra jaraw kemm huma attivi certi partijiet tal-moħħi u jistgħu jgħinu wkoll biex jagħżlu t-tip ta' dimenžja.
- Assessjar psikjatriku - biex ikunu identifikati l-kundizzjonijiet li jistgħu jiġu kukturati bħad-dipressjoni li tista' tkun tixxbiha lid-dimenzja u biex ikunu mmaniġjati s-sintomi psikjatriċi bhall-ansjetà jew id-delużjonijiet li jistgħu jkunu preżenti flimkien mad-dimenzja.

Fejn tibda

L-aħjar post fejn tibda huwa mat-tabib tal-pazjent. Wara li jistudja s-sintomi u jordna t-testijiet bl-iskrining, it-tabib ikun jista' jagħmel dijanjosi preliminary u idealment jirreferi l-persuna lejn speċjalista mediku bħal newrologu, ġerjatra jew psikjatra.

Xi wħud jistgħu ma jaqblux mal-idea li jmorru jaraw tabib. Xi drabi wħud ma jirrealizzawx li mhumiex f'sikkithom għax l-bidliet fil-moħħi li jsiru bid-dimenzja jfixxu l-kapacità li persuna tagħraf jew tapprezzza it-tibdiliet li jkunu qiegħdin isiru. Oħra jkollhom ħji tal-kundizzjoni tagħhom, jistgħu jkunu jibzgħu li jkollhom il-biza' tagħhom ikkonfermat. Wieħed mill-aħjar modi biex tingħelex din il-problema huwa li tinstab raġuni oħra għal żjara lit-tabib. Forsi tissuġġerixxi biex tiċċekkja l-pressjoni tad-demm jew reviżjoni ta' xi kura jew kundizzjon li tkun ilha sejra għal žmien twil. Mod ieħor huwa li tissuġġerixxi li t-tnejn tkunu wasaltu biex ikollkom eżami fiziku. Attitudni kalma f'dan iż-żmien tista' tgħin biex jingħelbu l-biza' u t-thassib tal-persuna.

Jekk il-persuna tibqa' ma tridx tmur għand it-tabib:

- Tkellem mat-tabib tal-persuna biex jagħtik parir
- Ikkuntattja lit-Tim tal-Assessor tal-Kura għall-Anzjani (Aged Care Assessment Team – ACAT) permezz ta' My Aged Care fuq **1800 200 422** għall-informazzjoni.
- Ċempel il-Linja Nazzjonali tal-Ġħajjnuna fid-Dimenzja fuq **1800 100 500**

Jekk tħoss li t-tabib tiegħek mhuwiex jieħu bis-serjetà it-thassib tiegħek ara tieħux opinjoni oħra.

Għal informazzjoni iktar dettaljata dwar is-sinjalib bikrin tad-dimenzja u kif titlob l-ġħajjnuna ta' tabib niżżejjel il-ktejjeb *Inkwetata Dwar il-Memoria Tiegħek?* Mill-websajt ta' Dementia Australia jew ċempel il-Linja Nazzjonali tal-Ġħajjnuna fid-Dimenzja biex tordna kopja.

AKTAR TAGħrif

Dementia Australia toffri rapport, tagħrif, edukazzjoni u pariri. Ikkuntattja l-Linja Nazzjonali ta' Ghajjnuna fid-Dimenzja fuq **1800 100 500** jew mur fuq il-websajt tagħha dementia.org.au



Għall-ġħajjnuna fil-lingwa ċempel is-Servizz Telefoniku tat-Traduzzjoni u l-Interpretar fuq **131 450**

Diagnosing dementia

This Help Sheet provides information about the early signs of dementia, the techniques used to diagnose dementia and the importance of an early and correct diagnosis.

What are the early signs of dementia?

Because the diseases that cause dementia develop gradually, the early signs may be very subtle and not immediately obvious. Early symptoms also depend on the type of dementia and vary a great deal from person to person.

Common early symptoms include:

- Memory problems, particularly remembering recent events
- Increasing confusion
- Reduced concentration
- Personality or behaviour changes
- Apathy and withdrawal or depression
- Loss of ability to do everyday tasks

Sometimes people fail to recognise that these symptoms indicate that something is wrong. They may mistakenly assume that such behaviour is a normal part of ageing, or symptoms may develop so gradually they go unnoticed for a long time. Sometimes people may be reluctant to act even when they know something is wrong.

For the person experiencing the symptoms, the very nature of these changes within the brain may mean that the person is unable to recognise that there are changes.

Warning signs

This is a checklist of common symptoms of dementia. Go through the list and tick any symptoms that are present. If there are several ticks, consult a doctor for a complete assessment.

Memory loss that affects day-to-day function

It's normal to occasionally forget appointments and remember them later. A person with dementia may forget things more often or not remember them at all.

Difficulty performing familiar tasks

People can get distracted and they may forget to serve part of a meal. A person with dementia may have trouble with all the steps involved in preparing a meal.

Disorientation to time and place

A person with dementia may have difficulty finding their way to a familiar place, or feel confused about where they are, or think they are back in some past time of their life.

Problems with language

Everyone has trouble finding the right word sometimes, but a person with dementia may forget simple words or substitute inappropriate words, making them difficult to understand. They might also have trouble understanding others.

Problems with abstract thinking

Managing finances can be difficult for anyone, but a person with dementia may have trouble knowing what the numbers mean or what to do with them.

Poor or decreased judgement

Many activities require good judgement. When this ability is affected by dementia, the person may have difficulty making appropriate decisions, such as what to wear in cold weather.

Problems with spatial skills

A person with dementia may have difficulty judging distance or direction when driving a car.

Problems misplacing things

Anyone can temporarily misplace a wallet or keys. A person with dementia may often put things in inappropriate places.

Changes in mood, personality or behaviour

Everyone becomes sad or moody from time to time. Someone with dementia can exhibit rapid mood swings for no apparent reason. They can become confused, suspicious or withdrawn. Some can become disinhibited or more outgoing.

A loss of initiative

It's normal to tire of some activities. However, dementia may cause a person to lose interest in previously enjoyed activities.

National Dementia Helpline **1800 100 500**

dementia.org.au

This help sheet is funded by the Australian Government

It may not be dementia

Remember that many conditions have symptoms similar to dementia, so do not assume that someone has dementia just because some of the above symptoms are present. Strokes, depression, alcoholism, infections, hormone disorders, nutritional deficiencies and brain tumours can all cause dementia-like symptoms. Many of these conditions can be treated.

A correct diagnosis is important

Consulting a doctor at an early stage is critical. Only a medical practitioner can diagnose dementia. A complete medical assessment may identify a treatable condition and ensure that it is treated correctly, or it might confirm the presence of dementia.

An assessment may include the following:

- Medical history – the doctor will ask about past and current medical problems, family medical history, any medications being taken, and the problems with memory, thinking or behaviour that are causing concern. The doctor may also wish to speak to a close family member who can help provide all the necessary information.
- Physical examination – this may include tests of the senses and movement function, as well as heart and lung function, to help rule out other conditions.
- Laboratory tests – will include a variety of blood and urine tests to identify any possible illness which could be responsible for the symptoms. In some cases, a small sample of spinal fluid may be collected for testing.
- Neuropsychological or cognitive testing – a variety of tests are used to assess thinking abilities including memory, language, attention and problem solving. This can help identify specific problem areas, which in turn helps identify the underlying cause or the type of dementia.
- Brain imaging – there are certain scans that look at the structure of the brain and are used to rule out brain tumours or blood clots in the brain as the reason for symptoms, and to detect patterns of brain tissue loss that can differentiate between different types of dementia. Other scans look at how active certain parts of the brain are and can also help discriminate the type of dementia.
- Psychiatric assessment – to identify treatable disorders such as depression, and to manage any psychiatric symptoms such as anxiety or delusions which may occur alongside dementia

Where to begin

The best place to start is with the person's doctor. After considering the symptoms and ordering screening tests, the doctor may offer a preliminary diagnosis and will, ideally, refer the person to a medical specialist such as a neurologist, geriatrician or psychiatrist.

Some people may be resistant to the idea of visiting a doctor. Sometimes people do not realise that there is anything wrong with them because the brain changes that occur with dementia interfere with the ability to recognise or appreciate the changes occurring. Others, who do have insight into their condition, may be afraid of having their fears confirmed. One of the best ways to overcome this problem is to find another reason to visit the doctor. Perhaps suggest a blood pressure check or a review of a long-term condition or medication. Another way is to suggest that it is time for both of you to have a physical check-up. A calm attitude at this time can help overcome the person's worries and fears.

If the person still will not visit the doctor:

- Talk to the person's doctor for advice
- Contact your local Aged Care Assessment Team (ACAT) via My Aged Care on **1800 200 422** for information
- Call the National Dementia Helpline on **1800 100 500**

If you feel your doctor is not taking your concerns seriously, consider seeking a second opinion.

For more detailed information about the early signs of dementia and seeking help from a doctor, download the **Worried About Your Memory?** booklet from the Dementia Australia website or call the National Dementia Helpline to order a copy.

FURTHER INFORMATION

Dementia Australia offers support, information, education and counselling. Contact the National Dementia Helpline on **1800 100 500**, or visit our website at dementia.org.au



For language assistance phone the
Translating and Interpreting Service
on **131 450**